

## Tilburg University

### Digraph competitions and cooperative games

van den Brink, J.R.; Borm, P.E.M.

*Publication date:*  
1994

[Link to publication in Tilburg University Research Portal](#)

*Citation for published version (APA):*

van den Brink, J. R., & Borm, P. E. M. (1994). *Digraph competitions and cooperative games*. (CentER Discussion Paper; Vol. 1994-24). Unknown Publisher.

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

#### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CBM

8414

1994 24

8414

1994

NR.24

entER


for

omic Research

# Discussion paper



\* C I N O 1 5 3 7 \*



Center  
for  
Economic Research

8417  
1004  
24

No. 9424

79

**DIGRAPH COMPETITIONS AND  
COOPERATIVE GAMES**

by René van den Brink  
and Peter Borm

February 1994

ISSN 0924-7815



# Digraph Competitions and Cooperative Games\*

René van den Brink<sup>†</sup>

Peter Borm

Department of Econometrics

Tilburg University

P.O. Box 90153

5000 LE Tilburg

The Netherlands

February 1994

---

\*The authors would like to thank Vincent Feltkamp, Rob Gilles and Stef Tijs for their useful remarks.

<sup>†</sup>This author is financially supported by the Netherlands Organization for Scientific Research (NWO), grant 450-228-022

## Abstract

A dominance structure or competition between a set of players can be modelled by means of a *directed graph*  $(N, D)$ . Here, the set of nodes  $N$  represents a set of players, and  $(i, j) \in D$  means that player  $i$  dominates or defeats player  $j$ . We introduce the *conservative score mapping* which assigns to every digraph a *cooperative game with transferable utilities*. Solution concepts for these score games can be interpreted as a way to evaluate the underlying dominance structure or competition, i.e., as a way to rank the players.

We provide characterizations of the score mapping and of the subclass of score games. It is shown that a score game is *convex*, and its *core* is equal to the convex hull of the set of *score vectors* corresponding to those subgraphs in which each player that is defeated in the original digraph now is defeated exactly once. Consequently, all *marginal vectors* are such score vectors. However, in general the converse need not hold, and the class of digraphs for which it does is characterized.

Furthermore, the *Shapley value* of a score game, which by definition is the mean of all marginal vectors, turns out to be also the mean of all score vectors.

## 1 Introduction

A situation in which a set of players can obtain certain payoffs by cooperation can be represented by a *cooperative game with transferable utilities* – or simply a TU-game – being a pair  $(N, v)$  where  $N$  denotes the set of players and the *characteristic function*  $v: 2^N \rightarrow \mathbb{R}$  is such that  $v(\emptyset) = 0$ . In the sequel we take the set  $N$  to be fixed and therefore represent each TU-game on  $N$  by its characteristic function. We denote the set of all TU-games on  $N$  by  $\mathcal{G}^N$ . If we model a cooperative situation in this way then the players only differ with respect to their contributions to the values of the various coalitions but further they are assumed to be socially symmetric. In recent literature models have been developed in which the players are part of a social organization structure which limits the possibilities of coalition formation. In some of these models this social structure is represented by a *graph*. As an example we mention the models in which there are limited communication possibilities between the players which are represented by an *undirected graph*. It is assumed that players can only cooperate if they can communicate with one another. For this we refer to Myerson

(1977), Owen (1986), Borm, Owen and Tijs (1992), and the survey paper by Borm, van den Nouweland and Tijs (1993). Another approach introducing social asymmetries between the players can be found in, e.g., Gilles, Owen and van den Brink (1992) and Faigle and Kern (1993). Here the players are part of some hierarchical organization which is represented by a *directed graph*. It is assumed that players can only cooperate if they have permission from their superiors in the hierarchy.

In all models mentioned above a directed or undirected graph limits the possibilities of coalition formation in a TU-game. From a TU-game and a graph another TU-game is derived taking into account the limited possibilities of cooperation. Of course, the position of a player in the graph affects his role in the modified game.

In this paper we will derive a TU-game from just a graph. Such a TU-game can be seen as a *social power measure* in the sense that the value of a coalition is a measure of the importance of the coalition in the social structure represented by the graph. We concentrate on directed graphs.

A directed graph or *digraph* is a pair  $(N, D)$  where  $N$  is a finite set of nodes and  $D \subset N \times N$  is a binary relation on  $N$ . Such a digraph can represent various hierarchical organizations. Above we already mentioned the interpretation as a permission structure in which there are players that need permission from their superiors before they can cooperate. A digraph also can be interpreted as an exchange economy such that the nodes represent the agents that participate in the economy and the fact that agent  $i$  dominates agent  $j$  means that agent  $i$  sets the conditions under which trade between agent  $j$  and himself will take place. (For example, he sets the prices under which he exchanges commodities with agent  $j$ .) This interpretation is considered in van den Brink and Gilles (1993). Finally we mention two closely related interpretations of digraphs that can be found in, e.g., Rubinstein (1980) and Laffont, Laslier and Le Breton (1993). One is the interpretation of a digraph as a *preference relation* such that the nodes represent alternatives between which an agent can choose. The fact that alternative  $i$  dominates alternative  $j$  then means that an agent or group of agents prefers  $i$  to  $j$  when comparing these alternatives pairwise. The other is the interpretation of a digraph as a *competition* between players or teams. Then the nodes represent the players or teams that participate in the competition and  $(i, j) \in D$  means



that player  $i$  has won the match he played against player  $j$ .

In the sequel we do not want to fix the specific context. The various results are formulated in a general setting. In section 2 we introduce and axiomatize a specific mapping that assigns a (convex) TU-game to every digraph on the set  $N$ . The games that can be derived in this way are called *conservative score games* and form a subclass of  $\mathcal{G}^N$ . We provide a characterization of this subclass. It is seen that a conservative score game is convex and in section 3 we show that the core of such a game coincides with the convex hull of the set of score vectors corresponding to those subgraphs of  $D$  in which each player that is dominated in  $D$  is dominated by exactly one other player. As a consequence all marginal vectors are such score vectors. The class of digraphs for which the converse also holds is characterized by means of the non-existence of so-called *anti-directed semi-circuits*. In section 4 we consider the Shapley value (Shapley (1953)) of a conservative score game and argue that it is equal to the *BG-measure* which is introduced in van den Brink and Gilles (1992) as a *power measure* for digraphs. It is shown that the Shapley value of a conservative score game, which by definition is the mean of all marginal vectors, is also the mean of the score vectors that are considered in section 3.

We conclude this section by presenting some concepts about digraphs that will be used in the sequel. We only consider *finite, irreflexive* digraphs. A digraph  $(N, D)$  is finite if  $N$  is finite and it is irreflexive if  $(i, i) \notin D$  for every  $i \in N$ . We simply refer to these graphs as digraphs. In the sequel we assume the set  $N$  to be fixed and therefore we represent a digraph by its binary relation. The collection of all digraphs on  $N$  is denoted by  $\mathcal{D}^N$ .

Let  $D \in \mathcal{D}^N$ . For  $E \subset N$ , the nodes in

$$S_D(E) := \{j \in N \mid \text{there is an } i \in E \text{ such that } (i, j) \in D\}$$

are called the *successors* of  $E$  in  $D$ . For node  $i \in N$  the nodes in

$$P_D(i) := \{j \in N \mid (j, i) \in D\}$$

are called the *predecessors* of node  $i$  in  $D$ . (We will often omit the subscript  $D$ .) The *score measure* on  $N$  is the function  $\sigma: \mathcal{D}^N \rightarrow \mathbb{R}^N$  that assigns to every node



$i$  in digraph  $D$  the number of successors of  $i$ . A digraph  $A$  is a *single predecessor digraph* if  $\#P_A(i) \leq 1$  for all  $i \in N$ . A single predecessor digraph in  $D \in \mathcal{D}^N$  is a single predecessor digraph  $A \subset D$  such that  $P_A(i) \neq \emptyset$  if and only if  $P_D(i) \neq \emptyset$ . The collection of all single predecessor digraphs in  $D$  is denoted by  $\mathcal{A}_D$ .

## 2 The conservative score game

In this section we introduce a specific mapping that assigns a cooperative game with transferable utilities to every digraph on the set of nodes  $N$ . The set of players in such a TU-game corresponds to the nodes of the digraph while the characteristic function is given in the following definition.

**Definition 2.1** *The conservative score mapping on  $N$  is the mapping  $v^c: \mathcal{D}^N \rightarrow \mathcal{G}^N$  given by*

$$v^c(D)(E) = \#\{j \in S(E) \mid P(j) \subset E\} \text{ for all } E \subset N \text{ and } D \in \mathcal{D}^N.$$

For  $D \in \mathcal{D}^N$  the game  $v^c(D)$  is called the *conservative score game* corresponding to  $D$  and it assigns to every coalition  $E \subset N$  the number of successors of  $E$  that have no predecessors outside  $E$ . If we interpret this number as a measure of the importance of coalition  $E$  in the digraph then the conservative score game can be seen as a *social power measure* that measures the ‘dominance power’ of coalitions of nodes in a digraph. Using *unanimity games*  $u_T$ , which for every  $T \in 2^N \setminus \{\emptyset\}$  are given by

$$u_T(E) = \begin{cases} 1 & \text{if } E \supset T \\ 0 & \text{else.} \end{cases}$$

it is easy to see that a conservative score game can be expressed as in the following lemma.

**Lemma 2.2** *For every  $D \in \mathcal{D}^N$  it holds that*

$$v^c(D) = \sum_{i \in S(N)} u_{P(i)}.$$

**Example 2.3** Consider the digraph  $D = \{(1, 2), (3, 2), (2, 4), (3, 4)\}$  on  $N = \{1, \dots, 4\}$ . Since  $P(1) = P(3) = \emptyset$ ,  $P(2) = \{1, 3\}$  and  $P(4) = \{2, 3\}$  it follows from Lemma 2.2 that  $v^c(D) = u_{\{1,3\}} + u_{\{2,3\}}$ .

From the fact that each conservative score game can be expressed as a positive sum of unanimity games it follows that the conservative score game corresponding to  $D \in \mathcal{D}^N$  is *convex* which means that

$$v^c(D)(E) + v^c(D)(F) \leq v^c(D)(E \cup F) + v^c(D)(E \cap F) \text{ for all } E, F \subset N.$$

Next we give four axioms on a mapping  $v: \mathcal{D}^N \rightarrow \mathcal{G}^N$  that uniquely determine the conservative score mapping. The first axiom states that the value of the grand coalition  $N$  is equal to the total number of dominated nodes in the digraph.

**Axiom 2.4 (Efficiency)** *For every  $D \in \mathcal{D}^N$  it holds that*

$$v(D)(N) = \#S(N).$$

The second axiom states that a node that does not dominate any other node in the digraph adds the value zero to every coalition in the corresponding game.

**Axiom 2.5 (Dummy property)** *For every  $D \in \mathcal{D}^N$  and  $i \in N$  with  $S(i) = \emptyset$  it holds that*

$$v(D)(E) = v(D)(E \setminus \{i\}) \text{ for all } E \subset N.$$

The third axiom states that a node that dominates all dominated nodes in the digraph is necessary for any coalition to obtain a non-zero payoff in the corresponding game.

**Axiom 2.6 (Top property)** *For every  $D \in \mathcal{D}^N$  and  $i \in N$  with  $S(i) = S(N)$  it holds that*

$$v(D)(E) = 0 \text{ for all } E \subset N \setminus \{i\}.$$

The fourth axiom states that if we consider the union of two or more digraphs on the same set of nodes which are such that each node is dominated in at most one of these digraphs then the corresponding game equals the sum of the games corresponding to the constituting subdigraphs. In order to formalize this idea we introduce the following definitions. A *partition* of  $D \in \mathcal{D}^N$  is a collection  $\mathcal{P} = \{D_1, \dots, D_t\}$  of digraphs such that the following two conditions are satisfied:

- $\bigcup_{k=1}^t D_k = D$ ;
- $D_k \cap D_l = \emptyset$  for all  $1 \leq k, l \leq t, k \neq l$ .

A partition  $\mathcal{P}$  of a digraph  $D$  is *independent* if each node is dominated in at most one digraph in this partition, i.e, if besides the two conditions stated above  $\mathcal{P}$  also satisfies:

- $\#\{A \in \mathcal{P} \mid P_A(i) \neq \emptyset\} \leq 1$  for all  $i \in N$ .

**Axiom 2.7 (Additivity over independent partitions)** *For every  $D \in \mathcal{D}^N$  and each independent partition  $\mathcal{P}$  of  $D$  it holds that*

$$v(D) = \sum_{A \in \mathcal{P}} v(A).$$

The four axioms introduced above uniquely determine the conservative score mapping.

**Theorem 2.8** *A mapping  $v: \mathcal{D}^N \rightarrow \mathcal{G}^N$  is equal to the conservative score mapping if and only if it satisfies efficiency, the dummy property, the top property and additivity over independent partitions.*

#### PROOF

It can easily be verified that the conservative score mapping satisfies the four axioms.

Now let  $v: \mathcal{D}^N \rightarrow \mathcal{G}^N$  satisfy the four axioms and let  $D \in \mathcal{D}^N$ .

For every  $j \in S(N)$  we introduce the digraph  $D_j \in \mathcal{D}^N$  where

$$D_j := \{(i, j) \in D \mid i \in P(j)\}.$$

From efficiency of  $v$  it follows that  $v(D_j)(N) = 1$ . Let  $E \subset N$ .

If  $E \supset P(j)$  then  $S_{D_j}(i) = \emptyset$  for every  $i \in N \setminus E$ . The dummy property then implies that  $v(D_j)(E) = v(D_j)(N) = 1$ .

Otherwise, if  $E \not\supset P(j)$ , then  $P(j) \cap (N \setminus E) \neq \emptyset$ . Since  $S_{D_j}(i) = \{j\} = S_{D_j}(N)$  for all  $i \in P(j) \cap (N \setminus E)$ , the top property implies that  $v(D_j)(E) = 0$ .

The collection  $\{D_j\}_{j \in S(N)}$  is an independent partition of  $D$ . From additivity over independent partitions it then follows that

$$v(D)(E) = \sum_{j \in S(N)} v(D_j)(E) = \#\{j \in S(E) \mid P(j) \subset E\} = v^c(D)(E) \text{ for all } E \subset N.$$

□

The conservative score mapping assigns a TU-game to every digraph on  $N$ . However, not every TU-game can be the conservative score game corresponding to some digraph. The following proposition provides a necessary and sufficient condition for a TU-game to be the conservative score game corresponding to some digraph.

**Proposition 2.9** *Let  $v \in \mathcal{G}^N$ . Then  $v$  is a conservative score game if and only if there exists a sequence  $\mathcal{T} = (T_1, \dots, T_t)$  of coalitions, and a sequence  $\mathcal{I} = (i_1, \dots, i_t)$  of players such that the following three conditions are satisfied:*

- (i)  $i_k \in N \setminus T_k$  for all  $k \in \{1, \dots, t\}$ ;
- (ii)  $i_k \neq i_l$  for all  $k, l \in \{1, \dots, t\}$ ,  $k \neq l$ ;
- (iii)  $v = \sum_{T \in \mathcal{T}} u_T$ .

**PROOF**

**Only if**

Suppose that  $v$  is the conservative score game corresponding to  $D \in \mathcal{D}^N$ . According to Lemma 2.2 it holds that  $v = \sum_{i \in S(N)} u_{P(i)}$ .

Let  $S(N) = \{s_1, \dots, s_t\}$ . Taking  $\mathcal{T} = (P(s_1), \dots, P(s_t))$  and  $\mathcal{I} = (s_1, \dots, s_t)$ , the conditions (i), (ii) and (iii) are satisfied.

**If**

Let  $\mathcal{T} = (T_1, \dots, T_t)$  and  $\mathcal{I} = (i_1, \dots, i_t)$  satisfy conditions (i), (ii) and (iii).

Let  $D \in \mathcal{D}^N$  be given by  $D = \bigcup_{k=1}^t D_k$  where  $D_k := \{(j, i_k) \mid j \in T_k\}$  for every  $k \in \{1, \dots, t\}$ . Then it readily follows that  $v$  is the conservative score game corresponding to  $D$ .

□

Thus Proposition 2.9 characterizes the class of all conservative score games on  $N$ . An alternative to the conservative score mapping is the *(optimistic) score mapping\** which is the mapping  $v^*: \mathcal{D}^N \rightarrow \mathcal{G}^N$  given by

$$v^*(D)(E) = \#S(E) \text{ for every } E \subset N \text{ and } D \in \mathcal{D}^N.$$

Thus the (optimistic) score game corresponding to  $D \in \mathcal{D}^N$  assigns to every coalition  $E \subset N$  the total number of successors of  $E$ , irrespective of the fact whether these successors have predecessors outside  $E$  or not. Thus this game can be seen as a generalization of the score measure that is given in section 1. For every  $D \in \mathcal{D}^N$  the corresponding (optimistic) score game is the *dual game* of the corresponding conservative score game.

**Lemma 2.10** *For every  $D \in \mathcal{D}^N$  it holds that*

$$v^c(D)(N) - v^c(D)(N \setminus E) = v^*(D)(E) \text{ for all } E \subset N.$$

**PROOF**

Let  $D \in \mathcal{D}^N$  and  $E \subset N$ . Then

$$\begin{aligned} v^c(D)(N) - v^c(D)(N \setminus E) &= \#\{j \in S(N) \mid P(j) \subset N\} \\ &\quad - \#\{j \in S(N \setminus E) \mid P(j) \subset N \setminus E\} \\ &= \#S(N) - \#\{j \in S(N) \mid P(j) \cap E = \emptyset\} \\ &= \#\{j \in S(N) \mid P(j) \cap E \neq \emptyset\} = \#S(E) = v^*(D)(E). \end{aligned}$$

---

\*The (optimistic) *score game* corresponding to a digraph is introduced in van den Brink and Gilles (1992).

□

The optimistic score mapping can be axiomatized by replacing the top property in Theorem 2.8 by a *dual top property* which states that for every  $D \in \mathcal{D}^N$  and  $i \in N$  with  $S(i) = S(N)$  it holds that  $v(D)(E) = v(D)(N)$  for all  $E \ni i$ .

**Example 2.11** In this example we illustrate the independence of the four axioms stated in Theorem 2.8.

1. Let the mapping  $v^1: \mathcal{D}^N \rightarrow \mathcal{G}^N$  be given by

$$v^1(D)(E) = \frac{1}{2} \cdot v^c(D)(E) \text{ for all } E \subset N \text{ and } D \in \mathcal{D}^N.$$

This mapping satisfies all four axioms except efficiency. For the digraph of Example 2.3 it holds that  $v^1(D)(N) = 1 < 2 = \#S(N)$ .

2. Let the mapping  $v^2: \mathcal{D}^N \rightarrow \mathcal{G}^N$  be given by

$$v^2(D)(E) = \#\{j \in S(E) \mid (P(j) \cup \{j\}) \subset E\} \text{ for all } E \subset N \text{ and } D \in \mathcal{D}^N.$$

This mapping satisfies all four axioms except the dummy property. For the digraph of Example 2.3 it holds that  $v^2(D)(\{2, 3\}) = 0$  and  $v^2(D)(\{2, 3, 4\}) = 1$  although  $S(4) = \emptyset$ .

3. Consider the score mapping  $v^s: \mathcal{D}^N \rightarrow \mathcal{G}^N$  that is discussed above. This mapping satisfies all four axioms except the top property. For the digraph of Example 2.3 it holds that  $v^s(D)(\{1\}) = 1$  although  $S(3) = \{2, 4\} = S(N)$ .
4. Let the mapping  $v^3: \mathcal{D}^N \rightarrow \mathcal{G}^N$  for every  $E \subset N$  and  $D \in \mathcal{D}^N$  be given by

$$v^3(D)(E) = \#\{j \in S(E) \cup S(S(E)) \mid P(j) \cup P(P(j)) \subset E\}.$$

This mapping satisfies all four axioms except additivity over independent partitions. The two digraphs  $D^1 = \{(1, 2), (3, 2)\}$  and  $D^2 = \{(2, 4), (3, 4)\}$  form an independent partition of the digraph  $D$  of Example 2.3. However,

$$v^3(D^1)(\{2, 3\}) + v^3(D^2)(\{2, 3\}) = 0 + 1 \neq 0 = v^3(D)(\{2, 3\}).$$



### 3 The core of a conservative score game

In this section we consider the *core* of a conservative score game. The core of an arbitrary TU-game  $v \in \mathcal{G}^N$  is given by

$$Core(v) = \left\{ x \in \mathbb{R}^N \left| \begin{array}{c} \sum_{i \in N} x_i = v(N) \\ \text{and} \\ \sum_{i \in E} x_i \geq v(E) \text{ for all } E \subset N \end{array} \right. \right\}.$$

As stated in the previous section each conservative score game is *convex*. In Shapley (1971) it is shown that the core of a convex game  $v$  corresponds to the convex hull of the *marginal vectors*,  $\{m^\pi(v)\}_{\pi \in \Pi(N)}$ , where  $\Pi(N)$  denotes the collection of all permutations on  $N$  and for every  $\pi \in \Pi(N)$  and  $v \in \mathcal{G}^N$ ,  $m^\pi(v) \in \mathbb{R}^N$  is given by:

$$m_i^\pi(v) = v(\{j \in N \mid \pi(j) \leq \pi(i)\}) - v(\{j \in N \mid \pi(j) < \pi(i)\}) \text{ for all } i \in N. \quad (1)$$

Using this result we can characterize the core of a conservative score game. For every  $D \in \mathcal{D}^N$  we define

$$M(D) = \{m^\pi(v^c(D)) \in \mathbb{R}^N \mid \pi \in \Pi(N)\},$$

being the collection of all marginal vectors of the conservative score game corresponding to  $D$ , and

$$\Sigma(D) = \{\{\sigma(A) \in \mathbb{R}^N \mid A \in \mathcal{A}_D\},$$

being the collection of score measures of all single predecessor digraphs in  $D$ .

**Theorem 3.1** *For every  $D \in \mathcal{D}^N$  it holds that*

$$Core(v^c(D)) = Conv(\Sigma(D)).$$

**PROOF**

Let  $D \in \mathcal{D}^N$ . The proof of the theorem consists of the following steps.

- (a) We first prove that  $M(D) \subset \Sigma(D)$  by showing that for every  $\pi \in \Pi(N)$  there is an  $A \in \mathcal{A}_D$  such that  $m^\pi(v^c(D)) = \sigma(A)$ . Therefore let  $\pi \in \Pi(N)$ . Then



$$m_i^\pi(v^c(D)) = \#\{j \in S_D(i) \mid \pi(h) \leq \pi(i) \text{ for all } h \in P_D(j)\} \text{ for all } i \in N.$$

Let  $A \in \mathcal{D}^N$  be given by  $A = \{(i, j) \in D \mid \pi(h) \leq \pi(i) \text{ for all } h \in P_D(j)\}$ .

Since  $A \subset D$  and  $P_A(j) = \{i \in P_D(j) \mid \pi(i) \geq \pi(h) \text{ for all } h \in P_D(j)\}$  it is easy to see that  $A \in \mathcal{A}_D$  and  $\sigma(A) = m^\pi(v^c(D))$ .

- (b) Next we prove that  $\Sigma(D) \subset \text{Core}(v^c(D))$  by showing that  $\sigma(A) \in \text{Core}(v^c(D))$  for all  $A \in \mathcal{A}_D$ . Therefore let  $A \in \mathcal{A}_D$ .

By definition of a single predecessor digraph it holds that

$$\sum_{i \in N} \sigma_i(A) = \#S_D(N) = v^c(D)(N).$$

Further,  $\sum_{i \in E} \sigma_i(A) = \#\{j \in S_D(E) \mid P_A(j) \cap E \neq \emptyset\} \geq \#\{j \in S_D(E) \mid P_D(j) \subset E\} = v^c(D)(E)$  for all  $E \subset N$ .

Thus  $\sigma(A) \in \text{Core}(v^c(D))$ .

Thus we have proved that

$$M(D) \subset \Sigma(D) \subset \text{Core}(v^c(D)). \quad (2)$$

Since  $v^c(D)$  is convex it holds that  $\text{Core}(v^c(D)) = \text{Conv}(M(D))$ . With (2) it then follows that  $\text{Core}(v^c(D)) = \text{Conv}(\Sigma(D))$ .

□

Note that the number of permutations of the players in  $N$  is equal to  $\#\Pi(N) = (\#N)!$ , while the number of single predecessor digraphs in digraph  $D$  is equal to  $\#\mathcal{A}_D = \prod_{i \in S(N)} \#P(i)$ . Therefore, for many digraphs the number of single predecessor digraphs will be less than the number of permutations of  $N$ . However, this does not hold for all digraphs as can easily be seen by considering the *complete* digraph  $D = N \times N$ .

**Example 3.2** Reconsider the digraph  $D$  and corresponding conservative score game  $v$  of Example 2.3.

This digraph has 4 single predecessor digraphs:




Figure 1: The core of  $v$  of Example 3.2

$$A_1 = \{(1, 2), (2, 4)\}, \quad A_2 = \{(1, 2), (3, 4)\}, \quad A_3 = \{(3, 2), (2, 4)\}, \quad A_4 = \{(3, 2), (3, 4)\}.$$

The corresponding score measures are:

$$\sigma(A_1) = (1, 1, 0, 0), \quad \sigma(A_2) = (1, 0, 1, 0), \quad \sigma(A_3) = (0, 1, 1, 0), \quad \sigma(A_4) = (0, 0, 2, 0).$$

The core of  $v$  is represented by the shaded area in Figure 1. Note that the vectors  $\sigma(A_i)$ ,  $A_i \in \mathcal{A}_D$ , are precisely the extreme points of the core.


According to relation (2) in the proof of Theorem 3.1 each extreme point of the core of a conservative score game corresponds to the score measure of a single predecessor digraph in the corresponding digraph. In Example 3.2 the converse also holds. However, this is not always the case as the following example shows.

**Example 3.3** Consider the digraph  $D = \{(1, 2), (1, 5), (2, 4), (2, 5), (3, 2), (3, 4)\}$  on  $N = \{1, \dots, 5\}$  as represented in Figure 2.

This digraph has 8 single predecessor digraphs which are given by

$$A_1 = \{(1, 2), (1, 5), (2, 4)\}, \quad A_2 = \{(1, 2), (1, 5), (3, 4)\}, \quad A_3 = \{(1, 2), (2, 5), (2, 4)\},$$

$$A_4 = \{(1, 2), (2, 5), (3, 4)\}, \quad A_5 = \{(3, 2), (1, 5), (2, 4)\}, \quad A_6 = \{(3, 2), (1, 5), (3, 4)\},$$

Figure 2: Digraph  $D$  of Example 3.3

$$A_7 = \{(3, 2), (2, 5), (2, 4)\}, \text{ and } A_8 = \{(3, 2), (2, 5), (3, 4)\},$$

with score measures

$$\sigma(A_1) = (2, 1, 0, 0, 0), \sigma(A_2) = (2, 0, 1, 0, 0), \sigma(A_3) = (1, 2, 0, 0, 0),$$

$$\sigma(A_4) = (1, 1, 1, 0, 0), \sigma(A_5) = (1, 1, 1, 0, 0), \sigma(A_6) = (1, 0, 2, 0, 0),$$


$$\sigma(A_7) = (0, 2, 1, 0, 0), \text{ and } \sigma(A_8) = (0, 1, 2, 0, 0).$$

According to Lemma 2.2 the corresponding conservative score game is given by  $v = u_{\{1,2\}} + u_{\{1,3\}} + u_{\{2,3\}}$ . Thus

$$M(D) = \{(2, 1, 0, 0, 0), (2, 0, 1, 0, 0), (1, 0, 2, 0, 0), (1, 2, 0, 0, 0), (0, 2, 1, 0, 0), (0, 1, 2, 0, 0)\}.$$

The core of  $v$  and the score measures of the single predecessor digraphs in  $D$  are illustrated in Figure 3. (The core corresponds to the shaded area in the figure.)

Consider the score measure  $\sigma(A_4) = \sigma(A_5) = (1, 1, 1, 0, 0)$ . This score measure does not correspond to a marginal vector of  $v$ . The intuition behind this is as follows. Each pair of the players 1, 2 and 3 has a different successor in common. Therefore in any permutation of the players the marginal contribution of the first of the three players 1, 2 or 3, to enter is equal to 0, of the second to enter is equal to 1, and of the third to enter is equal to 2. Thus there is no permutation in which the marginal contribution of each of these three players is equal to 1.

Figure 3: The core of  $v$  of Example 3.3

Next we will characterize the class of digraphs  $D$  for which the set  $\Sigma(D)$  of score measures of all single predecessor digraphs coincides with the set  $M(D)$  of marginal vectors of the corresponding conservative score game. For this we introduce the following concept.

**Definition 3.4** Let  $D \in \mathcal{D}^N$ . A sequence of nodes  $(i_1, j_1, i_2, j_2, \dots, i_t, j_t)$ ,  $t \geq 2$ , is an **anti-directed semi-circuit** in  $D$  if the following conditions are satisfied

- (i)  $i_k \neq i_l$  and  $j_k \neq j_l$  for all  $k, l \in \{1, \dots, t\}$  with  $k \neq l$ ;
- (ii)  $j_k \in S(i_k) \cap S(i_{k+1})$  for all  $k \in \{1, \dots, t-1\}$  and  $j_t \in S(i_t) \cap S(i_1)$ .

Thus an anti-directed semi-circuit in a digraph  $D$  is a cycle such that the neighbours of each node  $i$  in the cycle are either both successors or both predecessors of  $i$  and, moreover, each node appears at most once as a successor and at most once as a predecessor in this cycle. In the digraph of Example 3.3 the sequence  $(1, 2, 3, 4, 2, 5)$  is an anti-directed semi-circuit. In the digraph of Examples 2.3 and 3.2 there does not exist an anti-directed semi-circuit.

**Theorem 3.5** Let  $D \in \mathcal{D}^N$ . Then

$\Sigma(D) = M(D)$  if and only if there is no anti-directed semi-circuit in  $D$ .

The proof of this theorem can be found in the appendix and is more or less based on the reasoning as given in Example 3.3.

## 4 The Shapley value of a conservative score game

In this section we consider the Shapley value of a conservative score game. By definition the Shapley value is the function  $Sh: \mathcal{G}^N \rightarrow \mathbb{R}^N$  which assigns to each TU-game the average of the marginal vectors of that game. In particular, for every  $D \in \mathcal{D}^N$  it holds that

$$Sh_i(v^c(D)) = \frac{1}{\#M(D)} \sum_{m \in M(D)} m_i \text{ for all } i \in N. \quad (3)$$

A *power measure* for digraphs is a function  $f: \mathcal{D}^N \rightarrow \mathbb{R}^N$  that assigns a real number to each node of a digraph which can be seen as a measure of the ‘dominance power’ of that node in the digraph. The score measure can be seen as such a power measure. Another power measure, introduced in van den Brink and Gilles (1992), is the *BG-measure*  $\beta: \mathcal{D}^N \rightarrow \mathbb{R}^N$ , which is given by

$$\beta_i(D) = \sum_{j \in S(i)} \frac{1}{\#P(j)} \text{ for all } i \in N \text{ and } D \in \mathcal{D}^N. \quad (4)$$

In section 2 we argued that the conservative score mapping assigns to every digraph a TU-game which can be seen as a social power measure that measures the dominance power of coalitions of nodes in the underlying digraph. Since the Shapley value assigns real numbers to all players in a TU-game, the Shapley value as expressed in equation (3) can be seen as a power measure for digraph  $D$ . It turns out that the Shapley value of a conservative score game is equal to the BG-measure of the underlying digraph. Moreover, this Shapley value, which by convexity of the game is the barycenter of the core (Shapley (1971)), is equal to the average of the score vectors corresponding to the single predecessor digraphs in the digraph. This is a rather surprising result since Example 3.3 shows that this set of score vectors may contain the extreme points of the core as a strict subset.

**Theorem 4.1** *Let  $D \in \mathcal{D}^N$ . Then  $Sh(v^c(D)) = \beta(D)$  and*

$$Sh_i(v^c(D)) = \frac{1}{\#\mathcal{A}_D} \sum_{A \in \mathcal{A}_D} \sigma_i(A) \text{ for all } i \in N.$$

## PROOF

Let  $D \in \mathcal{D}^N$ . According to Lemma 2.10 the conservative score game  $v^c(D)$  is the dual of the score game  $v^s(D)$ . In van den Brink and Gilles (1992) it is shown that  $Sh(v^s(D)) = \beta(D)$ . Since the Shapley value of a TU-game is equal to the Shapley value of its dual game it holds that  $Sh(v^c(D)) = \beta(D)$ .

For every  $(i, j) \in D$  we define  $\mathcal{A}_D(i, j) := \{A \in \mathcal{A}_D \mid j \in S_A(i)\}$ , being the collection of single predecessor digraphs in  $D$  in which  $i$  is the (unique) predecessor of  $j$ .

Then  $\#\mathcal{A}_D(i, j) = \prod_{h \in S(N) \setminus \{i, j\}} \#P(h)$ . Since  $\#\mathcal{A}_D = \prod_{h \in S(N)} \#P(h)$  it then holds that<sup>†</sup>

$$\begin{aligned} \beta_i(D) &= \sum_{j \in S_D(i)} \frac{1}{\#P(j)} = \frac{1}{\#\mathcal{A}_D} \sum_{j \in S_D(i)} \#\mathcal{A}_D(i, j) = \\ &= \frac{1}{\#\mathcal{A}_D} \sum_{j \in S_D(i)} \sum_{A \in \mathcal{A}_D(i, j)} 1 = \frac{1}{\#\mathcal{A}_D} \sum_{A \in \mathcal{A}_D} \sum_{j \in S_A(i)} 1 = \frac{1}{\#\mathcal{A}_D} \sum_{A \in \mathcal{A}_D} \sigma_i(A). \end{aligned}$$

□

**Example 4.2** Consider the digraph  $D$  of Example 3.3. For this digraph it holds that  $Sh(v^c(D)) = \beta(D) = (1, 1, 1, 0, 0)$ . Taking the average of the marginal vectors over all permutations of the players yields

$$\frac{1}{120} \begin{pmatrix} (20(2, 1, 0, 0, 0) + 20(2, 0, 1, 0, 0) + 20(1, 0, 2, 0, 0)) \\ + 20(1, 2, 0, 0, 0) + 20(0, 2, 1, 0, 0) + 20(0, 1, 2, 0, 0) \end{pmatrix} = (1, 1, 1, 0, 0),$$

while taking the average of the score measures of all single predecessor digraphs in  $D$  yields

$$\frac{1}{8} \begin{pmatrix} (2, 1, 0, 0, 0) + (2, 0, 1, 0, 0) + (1, 0, 2, 0, 0) + (1, 1, 1, 0, 0) \\ + (1, 1, 1, 0, 0) + (1, 2, 0, 0, 0) + (0, 2, 1, 0, 0) + (0, 1, 2, 0, 0) \end{pmatrix} = (1, 1, 1, 0, 0).$$

---

<sup>†</sup>This part is similar to the proof that the BG-measure of an acyclic, quasi-strongly connected digraph  $D$  is equal to the average of the score measures over all single predecessor digraphs in  $D$  as shown in van den Brink and Gilles (1993).



There are various axiomatizations of the Shapley value for arbitrary TU-games. One of these axiomatizations uses the efficiency, dummy player, necessary player and additivity axioms. A function  $f: \mathcal{G}^N \rightarrow \mathbb{R}^N$  is *efficient* if  $\sum_{i \in N} f_i(v) = v(N)$  for all  $v \in \mathcal{G}^N$ . It satisfies the *dummy player* property if  $f_i(v) = 0$  for all  $i \in N$  and  $v \in \mathcal{G}^N$  with  $v(E) = v(E \setminus \{i\})$  for all  $E \subset N$ . It satisfies the *necessary player* property if for all *monotone* games<sup>†</sup>  $v$  and  $i \in N$  such that  $v(E) = 0$  for all  $E \subset N \setminus \{i\}$  it holds that  $f_i(v) \geq f_j(v)$  for all  $j \in N$ . It is *additive* if for all  $v, w \in \mathcal{G}^N$  it holds that  $f(v) + f(w) = f(v + w)$ , where  $(v + w)(E) = v(E) + w(E)$  for all  $E \subset N$ . In order to axiomatize the Shapley value restricted to the class of conservative score games we need to adapt the additivity axiom since for two conservative score games  $v$  and  $w$  the sum game  $(v + w)$  need not to be a conservative score game as the following example shows.

**Example 4.3** Let  $N = \{1, 2, 3\}$ ,  $v = u_{\{1\}} + u_{\{1,2\}}$ , and  $w = u_{\{1,3\}}$ . The (unique) digraphs underlying these conservative score games are

$$D^1 = \{(1, 2), (1, 3), (2, 3)\} \text{ and } D^2 = \{(1, 2), (3, 2)\}.$$

Using Proposition 2.9 it can be seen that no digraph can be constructed that underlies the sum game  $(v + w) = u_{\{1\}} + u_{\{1,2\}} + u_{\{1,3\}}$ .

We say that a function  $f: \mathcal{G}^N \rightarrow \mathbb{R}^N$  is additive on a specific subclass  $\mathcal{G}$  of  $\mathcal{G}^N$  if for every pair of games  $v, w \in \mathcal{G}$  such that the sum game  $(v + w)$  is also an element of  $\mathcal{G}$  it holds that  $f(v + w) = f(v) + f(w)$ . Now the Shapley value restricted to the class of conservative score games can be axiomatized by the efficiency, dummy and necessary player axioms and additivity restricted to the class of conservative score games. (This can be seen by noting that each unanimity game, except the unanimity game on  $N$ , is a conservative score game and each conservative score game can be expressed as in Lemma 2.2.)

As stated in Theorem 4.1, the BG-measure of a digraph  $D$  is equal to the Shapley value of the conservative score game  $v^c(D)$ . It is interesting to note that by combining the set

---

<sup>†</sup>A TU-game  $v$  is monotone if  $v(E) \leq v(F)$  for all  $E \subset F \subset N$ .



of axioms that characterize the conservative score mapping and the set of axioms that characterize the Shapley value restricted to the class of conservative score games one obtains a set of axioms that uniquely determine the BG-measure as a power measure  $f: \mathcal{D}^N \rightarrow \mathbb{R}^N$ . For example, the dummy node properties of the conservative score mapping and the Shapley value, respectively, state that a node that does not dominate any other node in a digraph is a dummy player in the corresponding game, and that the Shapley value of a dummy player in a game is equal to 0. Combining these two axioms yields the dummy node property of the BG-measure which states that the BG-measure of a node that does not dominate any other node is equal to 0. Similarly, combining the other three properties yield the corresponding properties for the BG-measure. As can be derived from van den Brink and Gilles (1992) these four properties characterize the BG-measure.

## Appendix

Before proving Theorem 3.5 we introduce the following.

An anti-directed semi-circuit  $(i_1, j_1, \dots, i_t, j_t)$  is **minimal** in  $D$  if there is no anti-directed semi-circuit  $(h_1, g_1, \dots, h_s, g_s)$  in  $D$  such that  $\{h_1, \dots, h_s\} \subset \{i_1, \dots, i_t\}$ .

### PROOF OF THEOREM 3.5

Let  $D \in \mathcal{D}^N$ . Since  $M(D) \subset \Sigma(D)$  (as shown in the proof of Theorem 3.1) we have to prove that  $\Sigma(D) \subset M(D)$  if and only if there is no anti-directed semi-circuit in  $D$ .

### Only if

Suppose there is an anti-directed semi-circuit in  $D$ . Then there is a minimal anti-directed semi-circuit  $(i_1, j_1, \dots, i_t, j_t)$  in  $D$ .

Let  $I = \{i_1, \dots, i_t\}$  and  $J = \{j_1, \dots, j_t\}$ .

We distinguish the following two cases with respect to  $\#I$ .

1. Suppose that  $\#I = 2$ , i.e.,  $I = \{i_1, i_2\}$ .

Then we can construct an  $A \in \mathcal{A}_D$  that satisfies the following conditions:

- (i)  $P_A(j_1) = i_1$  and  $P_A(j_2) = i_2$ ;

(ii)  $P_A(h) \subset P_D(h) \cap I$  for all  $h \in S(I) \setminus J$ .

Since  $j_1 \notin S(i_1) \setminus S(i_2)$  and  $j_2 \notin S(i_2) \setminus S(i_1)$  it holds that

$$\sigma_{i_1}(A) \geq \#[S(i_1) \setminus S(i_2)] + 1 \text{ and } \sigma_{i_2}(A) \geq \#[S(i_2) \setminus S(i_1)] + 1.$$

Let  $\pi \in \Pi(N)$ . We may assume without loss of generality. that  $\pi(i_1) < \pi(i_2)$ .

Then  $m_{i_1}^\pi(v^c(D)) \leq \#[S(i_1) \setminus S(i_2)] < \#[S(i_1) \setminus S(i_2)] + 1$ .

Hence, there is no  $\pi \in \Pi(N)$  such that  $\sigma(A) = m^\pi(v^c(D))$ .

2. Suppose that  $\#I > 2$ .

(a) Let  $h \in S(I) \setminus J$ . We first prove that  $\#[P(h) \cap I] = 1$ .

Since  $h \in S(I)$  it holds that  $\#[P(h) \cap I] \geq 1$ .

Suppose that  $\#[P(h) \cap I] > 1$ . Then there exist  $k, l \in \{1, \dots, t\}$  with  $k < l$  and  $\{i_k, i_l\} \subset P(h)$ . We may assume without loss of generality. that  $k = 1$ .

If  $l < t$  then  $(i_1, j_1, \dots, i_l, j_l)$  is an anti-directed semi-circuit in  $D$  with  $\{i_1, \dots, i_l\} \subset I$  and  $\{i_1, \dots, i_l\} \neq I$ .

If  $l = t$  then  $(i_1, j_1, i_t, j_t)$  is an anti-directed semi-circuit in  $D$  with  $\{i_1, i_t\} \subset I$  and  $\{i_1, i_t\} \neq I$  (since by assumption  $\#I > 2$ ).

Both cases are in contradiction with  $(i_1, j_1, \dots, i_t, j_t)$  being a minimal anti-directed semi-circuit in  $D$ .

(b) Using (a) we can construct an  $A \in \mathcal{A}_D$  that satisfies the following conditions:

(i)  $P_A(j_k) = \{i_k\}$  for all  $k \in \{1, \dots, t\}$ ;

(ii)  $P_A(h) = P_D(h) \cap I$  for all  $h \in S(I) \setminus J$ .

(c) Next we prove that  $\sigma(A) \notin M(D)$ .

Since  $S_A(i_k) = S_D(i_k) \setminus \{j_{k-1}\}$  for all  $k \in \{2, \dots, t\}$  and  $S_A(i_1) = S_D(i_1) \setminus \{j_t\}$ , it holds that

$$\sigma_i(A) = \#S_D(i) - 1 \text{ for all } i \in I. \quad (5)$$

Let  $\pi \in \Pi(N)$  and let  $i \in I$  be such that  $\pi(i) \leq \pi(h)$  for all  $h \in I$ . Then  $\{j \in S_D(i) \mid P_D(j) \subset \{h \in N \mid \pi(h) \leq \pi(i)\}\} \cap J = \emptyset$ .

But then

$$m_i^\pi(v^c(D)) = \#\{j \in S_D(i) \mid P_D(j) \subset \{h \in N \mid \pi(h) \leq \pi(i)\}\} \leq \#S_D(i) - 2.$$

Hence there is no  $\pi \in \Pi(N)$  such that  $\sigma(A) = m^\pi(v^c(D))$ .

Thus we have proved that if there is an anti-directed semi-circuit in  $D$  then  $\Sigma(D) \not\subset M(D)$ .

**If**

Suppose there is no anti-directed semi-circuit in  $D$ , and let  $A \in \mathcal{A}_D$ . We show that there is a  $\pi \in \Pi(N)$  such that  $\sigma(A) = m^\pi(v^c(D))$ .

(a) We recursively define the sets  $L_k \subset N$ ,  $k \in \mathbb{N} \cup \{0\}$ , as follows:

$$L_0 := \emptyset$$

and

$$L_k := \left\{ i \in N \setminus \bigcup_{l=0}^{k-1} L_l \mid \begin{array}{l} \text{For every } j \in S_A(i) \text{ it holds that} \\ P_D(j) \subset \bigcup_{l=0}^{k-1} L_l \cup \{i\} \end{array} \right\} \text{ for all } k \in \mathbb{N}.$$

(b) Let  $k \in \mathbb{N}$  be such that  $N \setminus \bigcup_{l=0}^{k-1} L_l \neq \emptyset$ . We prove that  $L_k \neq \emptyset$ .

On the contrary suppose that  $L_k = \emptyset$ . Let  $\widehat{N} := N \setminus \bigcup_{l=0}^{k-1} L_l$ . Then for every  $i \in \widehat{N}$  there is a  $j \in S_A(i)$  such that  $P_D(j) \not\subset \bigcup_{l=0}^{k-1} L_l \cup \{i\}$  and thus  $[P_D(j) \cap \widehat{N}] \setminus \{i\} \neq \emptyset$ .

Let  $\widehat{n} = \#\widehat{N}$ . We can construct a sequence  $(i_1, j_1, i_2, j_2, \dots)$  satisfying the following conditions:

- (i)  $i_1 \in \widehat{N}$ ;
- (ii)  $j_k \in S_A(i_k)$  and  $i_{k+1} \in [P_D(j_k) \cap \widehat{N}] \setminus \{i_k\}$  for every  $k \in \{1, \dots, \widehat{n}\}$ .

Since  $\widehat{N}$  is finite there are  $k, l \in \{1, \dots, \widehat{n} + 1\}$  with  $k < l$ ,  $i_k = i_l$  and all players in  $\{i_k, \dots, i_{l-1}\}$  different.

By construction it holds that  $j_r \neq j_s$  for all  $r, s \in \{k, \dots, l-1\}$ .

But then  $(i_k, j_k, \dots, i_{l-1}, j_{l-1})$  is an anti-directed semi-circuit in  $D$ .

(c) From (a) and (b) it follows that there is an  $m \in \mathbf{N}$  such that

- (i)  $L_k \neq \emptyset$  for all  $k \in \{1, \dots, m\}$ ;
- (ii)  $L_k \cap L_l = \emptyset$  for all  $k, l \in \{1, \dots, m\}$  with  $k \neq l$ ;
- (iii)  $\bigcup_{k=1}^m L_k = N$ ,

i.e., the sets  $L_1, \dots, L_m$  form a partition of  $N$  consisting of non-empty sets only.

(d) Let  $\pi \in \Pi(N)$  be such that if  $i \in L_k$  and  $j \in L_l$  with  $k < l$  then  $\pi(i) < \pi(j)$ . We show that  $m^\pi(v^c(D)) = \sigma(A)$ .

Let  $i \in L_k$  for some  $1 \leq k \leq m$ . From (a) it follows that

- (i) If  $j \in S_A(i)$  then  $P_D(j) \subset \bigcup_{l=0}^{k-1} L_l \cup \{i\}$ ;
- (ii) If  $j \in S_D(i) \setminus S_A(i)$  and  $S_A(h) = \{j\}$  then  $h \in \bigcup_{l=k+1}^m L_l$  and thus  $P_D(j) \not\subset \bigcup_{l=0}^{k-1} L_l \cup \{i\}$ .

But then

$$\begin{aligned} m_i^\pi(v^c(D)) &= \#\{j \in S_D(i) \mid \pi(h) \leq \pi(i) \text{ for all } h \in P_D(j)\} \\ &= \#\left\{j \in S_D(i) \mid P_D(j) \subset \bigcup_{l=0}^{k-1} L_l \cup \{i\}\right\} = \sigma_i(A). \end{aligned}$$

Thus we have proved that if there is no anti-directed semi-circuit in  $D$  then  $\Sigma(D) \subset M(D)$ .

□

## References

- BEHZAD, M., G. CHARTRAND, AND L. LESNIAK-FOSTER (1979), *Graphs and Digraphs*, Wadsworth, Belmont, California.
- BORM, P., VAN DEN NOUWELAND, A., AND S. TIJS (1993), "Cooperation and Communication Restrictions: A Survey", forthcoming in Gilles, R.P., and Ruys, P.H.M. (eds.): *Imperfections and Behaviour in Economic Organizations: Proceedings of the conference held in Tilburg from April 17-19, 1990*, North Holland, Amsterdam.
- BORM, P., OWEN, G., AND S. TIJS (1992), "On the Positions of Communication Situations," *SIAM Journal on Discrete Mathematics*, 5, 305-320.
- BRINK, R. VAN DEN, AND R.P. GILLES (1993), "An Axiomatic Social Power Index for Hierarchically Structured Populations of Economic Agents", forthcoming in Gilles, R.P., and Ruys, P.H.M. (eds.): *Imperfections and Behaviour in Economic Organizations: Proceedings of the conference held in Tilburg from April 17-19, 1990*, North Holland, Amsterdam.
- BRINK, R. VAN DEN, AND R.P. GILLES (1992), *Measuring Domination in Directed Graphs*, Research Memorandum FEW 565, Department of Econometrics, Tilburg University, Tilburg.
- FAIGLE, U., AND W. KERN (1993), "The Shapley Value for Cooperative Games under Precedence Constraints", *International Journal of Game Theory*, 21, 249-266.
- GILLES, R.P., OWEN, G. AND J.R. VAN DEN BRINK (1992), "Games with Permission Structures: the Conjunctive Approach", *International Journal of Game Theory*, 20, 277-293.
- LAFFONT G., J.F. LASLIER, AND M. LEBRETON (1993), "The Bipartisan Set of a Tournament Game", *Games and Economic Behavior*, 5, 182-201.
- MYERSON, R.B. (1977), "Graphs and Cooperation in Games," *Mathematics of Operations Research*, 2, 225-229.
- OWEN, G. (1986) "Values of Graph-Restricted Games," *SIAM Journal of Algebraic Discrete Methods*, 7, 210-220.

- RUBINSTEIN, A. (1980), "Ranking the Participants in a Tournament", *SIAM Journal of Applied Mathematics*, 38, 108-111.
- SHAPLEY, L.S. (1953) "A Value for  $n$ -Person Games", *Annals of Mathematics Studies* 28 (Contributions to the Theory of Games Vol.2) (eds. H.W. Kuhn and A.W. Tucker), Princeton University Press, 307-317.
- SHAPLEY, L.S. (1971) "Cores of Convex Games", *International Journal of Game Theory*, 1, 11-26.



**Discussion Paper Series, CentER, Tilburg University, The Netherlands:**

(For previous papers please consult previous discussion papers.)

No.	Author(s)	Title
9314	O.P. Attanasio and M. Browning	Consumption over the Life Cycle and over the Business Cycle
9315	F. C. Drost and B. J. M. Werker	A Note on Robinson's Test of Independence
9316	H. Hamers, P. Borm and S. Tijs	On Games Corresponding to Sequencing Situations with Ready Times
9317	W. Güth	On Ultimatum Bargaining Experiments - A Personal Review
9318	M.J.G. van Eijs	On the Determination of the Control Parameters of the Optimal Can-order Policy
9319	S. Hurkens	Multi-sided Pre-play Communication by Burning Money
9320	J.J.G. Lemmen and S.C.W. Eijffinger	The Quantity Approach to Financial Integration: The Feldstein-Horioka Criterion Revisited
9321	A.L. Bovenberg and S. Smulders	Environmental Quality and Pollution-saving Technological Change in a Two-sector Endogenous Growth Model
9322	K.-E. Wärneryd	The Will to Save Money: an Essay on Economic Psychology
9323	D. Talman, Y. Yamamoto and Z. Yang	The $(2^{n+m+1} - 2)$ -Ray Algorithm: A New Variable Dimension Simplicial Algorithm For Computing Economic Equilibria on $S^n \times R_+^m$
9324	H. Huizinga	The Financing and Taxation of U.S. Direct Investment Abroad
9325	S.C.W. Eijffinger and E. Schaling	Central Bank Independence: Theory and Evidence
9326	T.C. To	Infant Industry Protection with Learning-by-Doing
9327	J.P.J.F. Scheepens	Bankruptcy Litigation and Optimal Debt Contracts
9328	T.C. To	Tariffs, Rent Extraction and Manipulation of Competition
9329	F. de Jong, T. Nijman and A. Röell	A Comparison of the Cost of Trading French Shares on the Paris Bourse and on SEAQ International
9330	H. Huizinga	The Welfare Effects of Individual Retirement Accounts
9331	H. Huizinga	Time Preference and International Tax Competition



No.	Author(s)	Title
9332	V. Feltkamp, A. Koster, A. van den Nouweland, P. Borm and S. Tijs	Linear Production with Transport of Products, Resources and Technology
9333	B. Lauterbach and U. Ben-Zion	Panic Behavior and the Performance of Circuit Breakers: Empirical Evidence
9334	B. Melenberg and A. van Soest	Semi-parametric Estimation of the Sample Selection Model
9335	A.L. Bovenberg and F. van der Ploeg	Green Policies and Public Finance in a Small Open Economy
9336	E. Schaling	On the Economic Independence of the Central Bank and the Persistence of Inflation
9337	G.-J. Otten	Characterizations of a Game Theoretical Cost Allocation Method
9338	M. Gradstein	Provision of Public Goods With Incomplete Information: Decentralization vs. Central Planning
9339	W. Güth and H. Kliemt	Competition or Co-operation
9340	T.C. To	Export Subsidies and Oligopoly with Switching Costs
9341	A. Demirgüç-Kunt and H. Huizinga	Barriers to Portfolio Investments in Emerging Stock Markets
9342	G.J. Almekinders	Theories on the Scope for Foreign Exchange Market Intervention
9343	E.R. van Dam and W.H. Haemers	Eigenvalues and the Diameter of Graphs
9344	H. Carlsson and S. Dasgupta	Noise-Proof Equilibria in Signaling Games
9345	F. van der Ploeg and A.L. Bovenberg	Environmental Policy, Public Goods and the Marginal Cost of Public Funds
9346	J.P.C. Blanc and R.D. van der Mei	The Power-series Algorithm Applied to Polling Systems with a Dormant Server
9347	J.P.C. Blanc	Performance Analysis and Optimization with the Power-series Algorithm
9348	R.M.W.J. Beetsma and F. van der Ploeg	Intramarginal Interventions, Bands and the Pattern of EMS Exchange Rate Distributions
9349	A. Simonovits	Intercohort Heterogeneity and Optimal Social Insurance Systems
9350	R.C. Douven and J.C. Engwerda	Is There Room for Convergence in the E.C.?

No.	Author(s)	Title
9351	F. Vella and M. Verbeek	Estimating and Interpreting Models with Endogenous Treatment Effects: The Relationship Between Competing Estimators of the Union Impact on Wages
9352	C. Meghir and G. Weber	Intertemporal Non-separability or Borrowing Restrictions? A Disaggregate Analysis Using the US CEX Panel
9353	V. Feltkamp	Alternative Axiomatic Characterizations of the Shapley and Banzhaf Values
9354	R.J. de Groof and M.A. van Tuijl	Aspects of Goods Market Integration. A Two-Country-Two-Sector Analysis
9355	Z. Yang	A Simplicial Algorithm for Computing Robust Stationary Points of a Continuous Function on the Unit Simplex
9356	E. van Damme and S. Hurkens	Commitment Robust Equilibria and Endogenous Timing
9357	W. Güth and B. Peleg	On Ring Formation In Auctions
9358	V. Bhaskar	Neutral Stability In Asymmetric Evolutionary Games
9359	F. Vella and M. Verbeek	Estimating and Testing Simultaneous Equation Panel Data Models with Censored Endogenous Variables
9360	W.B. van den Hout and J.P.C. Blanc	The Power-Series Algorithm Extended to the <i>BMAP/PH/1</i> Queue
9361	R. Heuts and J. de Klein	An $(s, q)$ Inventory Model with Stochastic and Interrelated Lead Times
9362	K.-E. Wärneryd	A Closer Look at Economic Psychology
9363	P.J.-J. Herings	On the Connectedness of the Set of Constrained Equilibria
9364	P.J.-J. Herings	A Note on "Macroeconomic Policy in a Two-Party System as a Repeated Game"
9365	F. van der Ploeg and A. L. Bovenberg	Direct Crowding Out, Optimal Taxation and Pollution Abatement
9366	M. Pradhan	Sector Participation in Labour Supply Models: Preferences or Rationing?
9367	H.G. Bloemen and A. Kapteyn	The Estimation of Utility Consistent Labor Supply Models by Means of Simulated Scores
9368	M.R. Baye, D. Kovenock and C.G. de Vries	The Solution to the Tullock Rent-Seeking Game When $R > 2$ : Mixed-Strategy Equilibria and Mean Dissipation Rates
9369	T. van de Klundert and S. Smulders	The Welfare Consequences of Different Regimes of Oligopolistic Competition in a Growing Economy with Firm-Specific Knowledge

No.	Author(s)	Title
9370	G. van der Laan and D. Talman	Intersection Theorems on the Simplotope
9371	S. Muto	Alternating-Move Preplays and $vN - M$ Stable Sets in Two Person Strategic Form Games
9372	S. Muto	Voters' Power in Indirect Voting Systems with Political Parties: the Square Root Effect
9373	S. Smulders and R. Gradus	Pollution Abatement and Long-term Growth
9374	C. Fernandez, J. Osiewalski and M.F.J. Steel	Marginal Equivalence in $v$ -Spherical Models
9375	E. van Damme	Evolutionary Game Theory
9376	P.M. Kort	Pollution Control and the Dynamics of the Firm: the Effects of Market Based Instruments on Optimal Firm Investments
9377	A. L. Bovenberg and F. van der Ploeg	Optimal Taxation, Public Goods and Environmental Policy with Involuntary Unemployment
9378	F. Thuijsman, B. Peleg, M. Amitai & A. Shmida	Automata, Matching and Foraging Behavior of Bees
9379	A. Lejour and H. Verbon	Capital Mobility and Social Insurance in an Integrated Market
9380	C. Fernandez, J. Osiewalski and M. Steel	The Continuous Multivariate Location-Scale Model Revisited: A Tale of Robustness
9381	F. de Jong	Specification, Solution and Estimation of a Discrete Time Target Zone Model of EMS Exchange Rates
9401	J.P.C. Kleijnen and R.Y. Rubinstein	Monte Carlo Sampling and Variance Reduction Techniques
9402	F.C. Drost and B.J.M. Werker	Closing the Garch Gap: Continuous Time Garch Modeling
9403	A. Kapteyn	The Measurement of Household Cost Functions: Revealed Preference Versus Subjective Measures
9404	H.G. Bloemen	Job Search, Search Intensity and Labour Market Transitions: An Empirical Exercise
9405	P.W.J. De Bijl	Moral Hazard and Noisy Information Disclosure
9406	A. De Waegenaere	Redistribution of Risk Through Incomplete Markets with Trading Constraints

No.	Author(s)	Title
9407	A. van den Nouweland, P. Borm, W. van Golstein Brouwers, R. Groot Bruinderink, and S. Tijs	A Game Theoretic Approach to Problems in Telecommunication
9408	A.L. Bovenberg and F. van der Ploeg	Consequences of Environmental Tax Reform for Involuntary Unemployment and Welfare
9409	P. Smit	Arnoldi Type Methods for Eigenvalue Calculation: Theory and Experiments
9410	J. Eichberger and D. Kelsey	Non-additive Beliefs and Game Theory
9411	N. Dagan, R. Serrano and O. Volij	A Non-cooperative View of Consistent Bankruptcy Rules
9412	H. Bester and E. Petrakis	Coupons and Oligopolistic Price Discrimination
9413	G. Koop, J. Osiewalski and M.F.J. Steel	Bayesian Efficiency Analysis with a Flexible Form: The AIM Cost Function
9414	C. Kilby	World Bank-Borrower Relations and Project Supervision
9415	H. Bester	A Bargaining Model of Financial Intermediation
9416	J.J.G. Lemmen and S.C.W. Eijffinger	The Price Approach to Financial Integration: Decomposing European Money Market Interest Rate Differentials
9417	J. de la Horra and C. Fernandez	Sensitivity to Prior Independence via Farlie-Gumbel-Morgenstern Model
9418	D. Tolman and Z. Yang	A Simplicial Algorithm for Computing Proper Nash Equilibria of Finite Games
9419	H.J. Bierens	Nonparametric Cointegration Tests
9420	G. van der Laan, D. Talman and Z. Yang	Intersection Theorems on Polytopes
9421	R. van den Brink and R.P. Gilles	Ranking the Nodes in Directed and Weighted Directed Graphs
9422	A. van Soest	Youth Minimum Wage Rates: The Dutch Experience
9423	N. Dagan and O. Volij	Bilateral Comparisons and Consistent Fair Division Rules in the Context of Bankruptcy Problems
9424	R. van den Brink and P. Borm	Digraph Competitions and Cooperative Games

P.O. BOX 90155, 5000 LE HEBORG, THE NETHERLANDS

**Bibliotheek K. U. Brabant**



**17 000 01176826 5**